



Event-specific Method for the Quantification of Maize Line LY038 Using Real-time PCR

Protocol

6 October 2008

**Joint Research Centre
Institute for Health and Consumer Protection
Biotechnology & GMOs Unit**

Method development:


Monsanto Company

Method validated by:

Community Reference Laboratory for GM Food and Feed (CRL-GMFF)
Biotechnology & GMOs Unit


Drafted by

E. Grazioli




Report Verification Team

1) M. Querci




2) M. Ermolli



Scientific and technical approval

M. Mazzara



Compliance to CRL Quality System

S. Cordeil



Authorisation to publish

G. Van den Eede



Address of contact laboratory:

European Commission, Joint Research Centre
Institute for Health and Consumer Protection (IHCP)
Biotechnology and GMOs Unit – Community Reference Laboratory for GM Food and Feed
Via Fermi 2749, 21027 Ispra (VA) - Italy

Content

1. GENERAL INFORMATION AND SUMMARY OF THE METHODOLOGY	4
2. VALIDATION STATUS AND PERFORMANCE CHARACTERISTICS	4
2.1 GENERAL	4
2.2 COLLABORATIVE TRIAL.....	5
2.3 LIMIT OF DETECTION (LOD)	5
2.4 LIMIT OF QUANTIFICATION (LOQ)	5
2.5 MOLECULAR SPECIFICITY	5
3. PROCEDURE	6
3.1 GENERAL INSTRUCTIONS AND PRECAUTIONS.....	6
3.2 REAL-TIME PCR FOR QUANTITATIVE ANALYSIS OF MAIZE EVENT LY038.....	7
3.2.1 General	7
3.2.2 Calibration	7
3.2.3 Real-time PCR set-up	7
3.3 DATA ANALYSIS	9
3.4 CALCULATION OF RESULTS	9
4. MATERIALS	10
4.1 EQUIPMENT	10
4.2 REAGENTS.....	10
4.3 PRIMERS AND PROBES	10

1. General information and summary of the methodology

This protocol describes an event-specific real-time quantitative TaqMan[®] PCR procedure for the determination of the relative content of maize event LY038 DNA to total maize DNA in a sample.

The PCR assay was optimised for use in real-time PCR instruments for plastic reaction vessels.

Template DNA extracted by means of suitable methods should be tested for quality and quantity prior to use in PCR assay. Tests for the presence of PCR inhibitors (e.g. monitor run of diluted series, use of DNA spikes) are recommended.

For the specific detection of maize event LY038 DNA, a 111-bp fragment of the integration region of the construct inserted into the plant genome (located at the 5' insert-to-plant junction) is amplified using two specific primers. PCR products are measured during each cycle (real-time) by means of a target-specific oligonucleotide probe labelled with the fluorescent dye 6-FAM as a reporter at its 5' end and with TAMRA as a quencher dye at its 3' end. The 5' nuclease activity of the Taq DNA polymerase results in the specific cleavage of the probe, leading to increased fluorescence, which is then monitored.

For the relative quantification of maize event LY038 DNA, a maize-specific reference system amplifies a 79-bp fragment of the maize endogenous *hmg* gene, using two specific primers and a *hmg* gene-specific probe labelled with 6-FAM and TAMRA as described above.

The measured fluorescence signal passes a threshold value after a certain number of cycles. This threshold cycle is called the "Ct" value. For quantification of the amount of event LY038 DNA in a test sample, LY038 and *hmg* Ct values are determined for the sample. Standard curves are then used to estimate the relative amount of maize event LY038 DNA to total maize DNA.

2. Validation status and performance characteristics

2.1 General

The method was optimised for suitable DNA extracted from maize seeds and grains containing mixtures of genetically modified and conventional maize.

The reproducibility and trueness of the method were tested through an international collaborative ring trial using DNA samples at different GMO contents.

2.2 Collaborative trial

The method was validated in an international collaborative study by the Joint Research Centre (JRC) of the European Commission. The study was undertaken with twelve participating laboratories in April/May 2007.

Each participant received twenty blind samples containing LY038 genomic DNA at five GM contents, ranging from 0.09% to 8.0%.

Each test sample was analysed by PCR in three repetitions. The study was designed as a blind quadruplicate collaborative trial; each laboratory received each level of GM LY038 in four unknown samples. Two replicates of each GM level were analysed on the same PCR plate.

A detailed validation report can be found at <http://gmo-crl.jrc.it/statusofdoss.htm>

2.3 Limit of detection (LOD)

According to the method developer, the relative LOD of the method is at least 0.045% in 200 ng of total Maize DNA. The relative LOD was not assessed in the collaborative study.

2.4 Limit of quantification (LOQ)

According to the method developer, the relative LOQ of the method is at least 0.09% in 200 ng of total Maize DNA. The lowest relative GM content of the target sequence included in collaborative trial was 0.09%.

2.5 Molecular specificity

According to the method developer, the method exploits a unique DNA sequence in the region of recombination between the insert and the plant genome. The sequence is specific to maize event LY038 and thus imparts event-specificity to the method.

The specificity of event-specific assay was experimentally tested by the applicant in real-time PCR against DNA extracted from plant materials containing the specific targets of Roundup Ready[®] corn MON88017, Roundup Ready[®] canola (RT200), Roundup Ready[®] canola (RT73), conventional canola, Roundup Ready[®] corn (GA21), Roundup Ready[®] maize (NK603), YieldGard[®] corn borer corn (MON810), YieldGard[®] rootworm corn (MON863), lysine maize (LY038), conventional corn, Roundup Ready[®] cotton (MON1445), Bollgard[®] cotton (MON531), Bollgard[®] cotton (MON757), BollgardII[®] cotton (MON15985), Roundup Ready[®] Flex cotton (MON 88913), conventional cotton, Roundup Ready[®] soybean (40-3-2), conventional soybean, Roundup Ready[®] wheat (MON71800), conventional wheat, lentil, sunflower, buckwheat, rye berries and peanut.

None of the GM materials tested, except the positive control maize line LY038, yielded detectable amplicons.

The specificity of the maize reference assay *hmg* was experimentally tested by the applicant against DNA extracted from plant materials containing Roundup Ready® corn MON88017, Roundup Ready® canola (RT200), Roundup Ready® canola (RT73), conventional canola, Roundup Ready® corn (GA21), Roundup Ready® corn (NK603), YieldGard® corn borer corn (MON810), YieldGard® rootworm corn (MON863), lysine maize (LY038), conventional corn, Roundup Ready® cotton (MON1445), Bollgard® cotton (MON531), Bollgard® cotton (MON757), BollgardII® cotton (MON15985), Roundup Ready® Flex cotton (MON 88913), conventional cotton, Roundup Ready® soybean (40-3-2), conventional soybean, Roundup Ready® wheat (MON71800), conventional wheat, lentil, sunflower, buckwheat, rye berries and peanut.

According to the applicant, only the positive control maize line LY038, Roundup Ready® corn (GA21), Roundup Ready® corn (NK603), YieldGard® corn borer corn (MON810), YieldGard® rootworm corn (MON863), Roundup Ready® corn MON88017 and conventional corn yielded detectable amplicons.

3. Procedure

3.1 General instructions and precautions

- The procedures require experience of working in sterile conditions.
- Laboratory organization, e.g. “forward flow direction” during PCR-setup, should follow international guidelines, e.g. ISO 24276:2006.
- PCR-reagents should be stored and handled in a separate room where no nucleic acids (with exception of PCR primers or probes) or DNA degrading or modifying enzymes have been handled previously. All handling of PCR reagents and controls requires dedicated equipment, especially pipettes.
- The equipment used should be sterilised prior to use and any residue of DNA has to be removed. All material used (e.g. vials, containers, pipette tips, etc.) must be suitable for PCR and molecular biology applications. They must be DNase-free, DNA-free, sterile and unable to adsorb protein or DNA.
- Filter pipette tips protected against aerosol should be used.
- Powder-free gloves should be used and changed frequently.
- Laboratory benches and equipment should be cleaned periodically with e.g. 10% sodium hypochloride solution (bleach).
- Pipettes should be checked regularly for precision and calibrated, if necessary.
- All handling steps - unless specified otherwise - should be carried out at 0 - 4°C.
- In order to avoid repeated freeze/thaw cycles aliquots should be prepared.

3.2 Real-time PCR for quantitative analysis of Maize event LY038

3.2.1 General

The PCR set-up for the taxon specific target sequence (*hmg*) and for the GMO (event LY038) target sequence should be carried out in separate vials. Multiplex PCR (using differential fluorescent labels for the probes) has not been tested or validated.

The use of maximum 200 ng of template DNA per reaction well is recommended.

The method is developed, for LY038, in a total volume of 50 µl per reaction mixture with the reagents as listed in Table 1 and, for *hmg*, in a total volume of 25 µl per reaction mixture with the reagents as listed in and Table 2.

3.2.2 Calibration

The calibration curves consist of five samples. The first point of the calibration curves is a 10% LY038 in non-GM maize DNA for a total of 200 ng of DNA (corresponding to approximately 73394 maize genome copies with one genome assumed to correspond to 2.725 pg of haploid maize genomic DNA) ⁽¹⁾.

A calibration curve is produced by plotting the Ct-values against the logarithm of the target copy number for the calibration points. This can be done e.g. by means of spreadsheet software, e.g. Microsoft Excel, or directly by options available with the sequence detection system software.

The copy number measured for the unknown sample DNA is obtained by interpolation from the standard curves.

3.2.3 Real-time PCR set-up

1. Thaw, mix gently and centrifuge the required amount of components needed for the run. Keep thawed reagents on ice.
2. In two reaction tubes (one for the LY038 system and one for the *hmg* system) on ice, add the following components (Tables 1 and 2) in the order mentioned below (except DNA) to prepare the master mixes.

Table 1. Amplification reaction mixture in the final volume/concentration per reaction well for the LY038 specific system.

Component	Final concentration	$\mu\text{L}/\text{reaction}$
TaqMan [®] Universal PCR Master Mix (2x)	1x	25
LY038 AF (10 μM)	150 nM	0.75
LY038 AR (10 μM)	150 nM	0.75
LY038 AP (5 μM)	50 nM	0.50
Nuclease free water	#	19
Template DNA (max 200 ng)	#	4.0
Total reaction volume:		50

Table 2. Amplification reaction mixture in the final volume/concentration per reaction well for the maize *hmg* reference system.

Component	Final concentration	$\mu\text{L}/\text{reaction}$
Buffer A (10x)	1x	2.5
<i>hmg</i> F (10 μM)	300 nM	0.75
<i>hmg</i> R (10 μM)	300 nM	0.75
<i>hmg</i> P (5 μM)	160 nM	0.80
MgCl ₂ (25 mM)	6.5 mM	6.5
dNTPs mix (10 mM)	200 μM	0.5
Nuclease free water	#	8.95
Template DNA (max 200 ng)	#	4.0
Total reaction volume:		25

3. Mix gently and centrifuge briefly.
4. Prepare two reaction tubes (one for the LY038 and one for the *hmg* master mixes) for each DNA sample to be tested (standard curve samples, unknown samples and control samples).
5. Add to each reaction tube the correct amount of master mix (e.g. $46 \times 3 = 138 \mu\text{L}$ master mix for three PCR repetitions for LY038 and $21 \times 3 = 63 \mu\text{L}$ master mix for three PCR repetitions for *hmg*). Add to each tube the correct amount of DNA (e.g. $4 \times 3 = 12 \mu\text{L}$ DNA for three PCR repetitions). Vortex each tubes for approx. 10 sec. This step is mandatory to reduce the variability among the repetitions of each sample to a minimum.
6. Spin down the tubes in a microcentrifuge. Aliquot 50 μL in each well for LY038 and 25 μL for *hmg*. Seal the reaction plate with optical cover or optical caps. Centrifuge the plate at low speed (e.g. approximately $250 \times g$ for 1 minute at 4°C to room temperature) to spin down the reaction mixture.
7. Place the plate into the instrument.
8. Run the PCR with cycling conditions described in Table 3:

Table 3. Cycling program for LY038 system and maize *hmg* system

Step	Stage	T°C	Time (sec)	Acquisition	Cycles	
1	UNG	50°C	120	No	1	
2	Initial denaturation	95°C	600	No	1	
3	Amplification	Denaturation	95°C	15	No	45
		Annealing & Extension	60°C	60	Yes	

3.3 Data analysis

After the real-time PCR, analyse the run following the procedure below:

- a) Set the threshold: display the amplification curves of one system (e.g. LY038) in logarithmic mode. Locate the threshold line in the area where the amplification profiles are parallel (exponential phase of PCR) and where there is no "fork effect" between repetitions of the same sample. Press the "update" button to ensure changes affect Ct values. Switch to the linear view mode by clicking on the Y axis of the amplification plot, and check that the threshold previously set falls within the geometric phase of the curves.
- b) Set the baseline: determine the cycle number at which the threshold line crosses the first amplification curve and set the baseline three cycles before that value (e.g. earliest Ct = 25, set the baseline crossing at Ct = 25 – 3 = 22).
- c) Save the settings.
- d) Repeat the procedure described in a) and b) on the amplification plots of the other system (e.g. *hmg* system).
- e) Save the settings and export all the data to a text file for further calculations.

3.4 Calculation of results

After having defined a threshold value within the logarithmic phase of amplification as described above, the instrument's software calculates the Ct-values for each reaction.

The standard curves are generated both for the *hmg* and the LY038 specific systems by plotting the Ct values measured for the calibration points against the logarithm of the DNA copy numbers and by fitting a linear regression line into these data.

Thereafter, the standard curves are used to estimate the copy numbers in the unknown sample DNA.

For the determination of the amount of event LY038 DNA in the unknown sample, the LY038 copy number is divided by the copy number of the Maize reference gene (*hmg*) and multiplied by 100 to obtain the percentage value ($GM\% = LY038/hmg \times 100$).

4. Materials

4.1 Equipment

- Real-time PCR instrument for plastic reaction vessels (glass capillaries are not recommended for the described buffer composition)
- Plastic reaction vessels suitable for real-time PCR instrument (enabling undisturbed fluorescence detection)
- Software for run analysis
- Microcentrifuge
- Micropipettes
- Vortex
- Rack for reaction tubes
- 1.5/2.0 mL reaction tubes

4.2 Reagents

- TaqMan[®] Universal PCR Master Mix (2X). Applied Biosystems Cat. No. 4304437
- 0.5 M EDTA. Sigma Cat. No. E-7647-01-0
- PCR Nucleotide Mix (10 mM dNTPs). Promega Cat. No. C114G
- TaqMan[®] 1000X Rxn Gold/Buffer A Pack. Applied Biosystems Cat. No. 4304441
- AmpliTaq Gold Polymerase. Applied Biosystems Cat. No. N808-0244
- Nuclease-free water. Sigma Cat. No. W-4502
- 1 M Tris-HCl, pH 8.0. Sigma Cat. No. T-3038

4.3 Primers and Probes

Name	Oligonucleotide DNA Sequence (5' to 3')
<i>LY038 target sequence</i>	
LY038 primer AF	5' – TGG GTT CAG TCT GCG AAT GTT – 3'
LY038 primer AR	5' – AGG AAT TCG ATA TCA AGC TTA TCG A – 3'
LY038 probe AP	6 – FAM - CGA GCG GAG TTT ATG GGT CGA CGG - TAMRA
<i>Reference gene hmg target sequence</i>	
<i>hmg</i> primer F	5' – TTG GAC TAG AAA TCT CGT GCT GA – 3'
<i>hmg</i> primer R	5' – GCT ACA TAG GGA GCC TTG TCC T – 3'
<i>hmg</i> probe P	6 – FAM - CAA TCC ACA CAA ACG CAC GCG TA - TAMRA